arXiv:1406.3606v1 [math.CV] 9 Jun 2014

DECAY RATES FOR CUSP FUNCTIONS

TRISTRAM DE PIRO

Abstract. We make some observations on the decay rates of the Fourier coefficients of cusp functions.

Definition 0.1. *We let $H(0,1)$ denote the restrictions of holomorphic functions on the open disc $D(0,1+\epsilon)$, for some $\epsilon > 0$, and $C^\infty(S^1)$, the set of smooth, complex-valued functions on the unit circle S^1. For $f \in H(0;1)$, with power series expansion $f(z) = \sum_{n \in \mathcal{Z}_{\geq 0}} a_n z^n$, we let $\overline{f}(n) = a_n$, for $n \in \mathcal{Z}_{\geq 0}$, and $f_1 = f|_{\partial D(0;1)}$. For $g \in C^\infty(S^1)$, and $n \in \mathcal{Z}$, we let;*

$$\hat{g}(n) = \frac{1}{2\pi} \int_0^{2\pi} g(\theta) e^{-in\theta} d\theta$$

denote the n'th Fourier coefficient.

Lemma 0.2. *With notation as in 0.1, if $f \in H(0,1)$, and $n \in \mathcal{Z}_{\geq 0}$, we have $\overline{f}(n) = \hat{f}_1(n)$.*

Proof. By the definition of residues for holomorphic function, Cauchy's residue theorem, see [2], and Fourier series, see [3], we have, for $n \in \mathcal{Z}_{\geq 0}$, that;

$$\overline{f}(n) = \frac{1}{2\pi i} \int_{\partial D(0,1)} \frac{f(z)}{z^{n+1}} dz$$

$$= \frac{1}{2\pi i} \int_0^{2\pi} \frac{f(e^{i\theta})}{e^{i(n+1)\theta}} i e^{i\theta} d\theta$$

$$= \frac{1}{2\pi} \int_0^{2\pi} \frac{f(e^{i\theta})}{e^{in\theta}} d\theta = \hat{f}_1(n)$$

□

Remarks 0.3. *By rescaling, we obtain a similar result for $0 < \delta < 1$ and $f \in H(0, 1-\delta)$, $f_{1-\delta} \in C^\infty(S^1)$, where $f_{1-\delta}(z) = f((1-\delta)z)$, for $|z| = 1$. Namely;*

$$\overline{f}(n) = \frac{1}{(1-\delta)^n} \hat{f}_{1-\delta}(n).$$

Definition 0.4. *We let $\mathcal{H}_+ = \{z \in \mathcal{C} : Im(z) > 0\}$ denote the upper half plane and $H(\mathcal{H}_+)$ denote the set of holomorphic functions on $\mathcal{H}_+$. We let $Cusp(\mathcal{H}_+) \subset H(\mathcal{H}_+)$ denote holomorphic functions on $\mathcal{H}_+$, satisfying the additional symmetry condition;*

(i). $g(z+1) = g(z)$, for $z \in \mathcal{C}$.

and the cusp condition;

(ii). $lim_{Im(z)\to\infty} g(z) = 0$ (we assume the limit is uniform in $Re(z)$)

Remarks 0.5. *Observe that cusp forms , see* [1]*, are special examples of cusp functions. The map $\Phi = exp(2\pi iz) : \mathcal{H}_+ \to D(0,1)$ is holomorphic, and, taking a principal branch of the logarithm $\Gamma = \frac{log(z)}{2\pi i} : (D(0,1) \setminus [0,1)) \to (\mathcal{H}_+ \cap U)$, where $U = \{z \in \mathcal{H}_+ : 0 < z < 1\}$, we obtain a holomorphic function $\Gamma^*(g)$ on $(D(0,1) \setminus [0,1))$. The condition (i) ensures that $\Gamma^*(g)$ extends uniquely to a holomorphic function f_0 on the annulus $D(0,1) \setminus \{0\}$, such that $\Phi^*(f_0) = g$. Using a Laurent's Theorem and the condition (ii), f_0 extends uniquely to a holomorphic function f on $D(0,1)$ with $f(0) = 0$. Taking the power series expansion $f(z) = \sum_{n\in\mathcal{Z}_{\geq 0}} a_n z^n$ on $D(0,1)$, we obtain a convergent expansion $g(z) = \sum_{n\in\mathcal{Z}_{\geq 0}} a_n exp(2\pi inz)$ on $\mathcal{H}_+$. Similarly to before, for $n \in \mathcal{Z}_{\geq 0}$, we let $\overline{g}(n) = a_n$.*

Lemma 0.6. *If $g \in Cusp(\mathcal{H}_+)$, then, for any $m \in \mathcal{Z}_{>0}$, there exist constants $C_m \in \mathcal{R}_{\geq 0}$ and $D_m \in \mathcal{Z}_{>0}$, such that;*

$|\overline{g}(n)| \leq C_m n^{-m}$, for all $n \geq D_m$

Proof. Let f denote the holomorphic function on $D(0,1)$, obtained in Remark 0.5. For any $0 < \delta < 1$, we have that $f \in H(0, 1-\delta)$. By Remarks 0.3, we have that $\overline{f}(n) = \frac{1}{(1-\delta)^n}\hat{f}_{1-\delta}(n)$, $(*)$. As $f_{1-\delta} \in C^\infty(S^1)$, by a simple extension of Corollary 2.4 in [3], we have that, for any $m \in \mathcal{Z}_{>0}$, there exist constants $C_m \in \mathcal{R}_{\geq 0}$ and $B_m \in \mathcal{Z}_{>0}$, such that;

$|\hat{f}_{1-\delta}(n)| \leq A_m n^{-m}$, for all $n \geq B_m$

Then, by $(*)$ and Remarks 0.5, we have;

$|\overline{g}(n)| = |\overline{f}(n)| = \frac{1}{(1-\delta)^n}|\hat{f}_{1-\delta}(n)| \leq C_m n^{-m}$, for all $n \geq B_m$

where $C_m = A_m n^{-m}$.

□

Remarks 0.7. *This result seems to improve significantly on the result of the Ramunajan-Petersson conjecture, that, for cusp forms of weight* $k \geq 2$, $|\overline{g}(n)| = O(n^{\frac{k-1}{2}+\gamma})$, $\gamma > 0$, $n \in \mathcal{Z}_{\geq 0}$. *The result also easily generalises to holomorphic functions on* $\mathcal{H}_+$, *satisfying just the symmetry condition* (i), *in Definition 0.4, for* $n \in \mathcal{Z}$, *with* $|n| \geq B_m$.

References

[1] Etale Cohomology and the Weil Conjecture, Eberhard Freitag, Springer-Verlag, Reinhardt Kiehl, (1987).

[2] Introduction to Complex Analysis, OUP, H.A. Priestley, (1990).

[3] Fourier Analysis, An Introduction, Elias Stein and Rami Shakarchi, Princeton Lectures in Analysis, (2003).

Mathematics Department, The University of Exeter, Exeter
E-mail address: tdpd201@exeter.ac.uk